

ARG20665 Goat IgG Isotype Control antibody (low endotoxin)

Package: 100 µg
Store at: -20°C

Summary

| | |
|---------------------|---|
| Product Description | Azide free and low endotoxin Goat Polyclonal antibody as a negative control antibody for Goat IgG |
| Tested Application | BL, ELISA, FACS, FLISA, IHC-Fr, IHC-P |
| Host | Goat |
| Clonality | Polyclonal |
| Isotype | IgG |
| Target Name | Goat IgG |
| Conjugation | Un-conjugated |

Application Instructions

| Application table | Application | Dilution |
|-------------------|--|-----------------|
| | BL | Assay-dependent |
| | ELISA | Assay-dependent |
| | FACS | Assay-dependent |
| | FLISA | Assay-dependent |
| | IHC-Fr | Assay-dependent |
| | IHC-P | Assay-dependent |
| Application Note | * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist. | |

Properties

| | |
|---------------------|--|
| Form | Liquid |
| Purification | Purified |
| Purification Note | Low endotoxin |
| Buffer | PBS |
| Concentration | 0.5 mg/ml |
| Storage instruction | For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use. |
| Note | For laboratory research only, not for drug, diagnostic or other use. |