

## ARG21408 anti-CD95 / Fas antibody [DX2] (Biotin)

Package: 50 tests

Store at: 4°C

### Summary

Product Description	Biotin-conjugated Mouse Monoclonal antibody [DX2] recognizes CD95 / Fas
Tested Reactivity	Hu, AGMK, R. Mk
Tested Application	ELISA, FACS, ICC/IF, IHC-Fr, IHC-P
Specificity	Human CD95.
Host	Mouse
Clonality	Monoclonal
Clone	DX2
Isotype	IgG1, kappa
Target Name	CD95 / Fas
Species	Human
Immunogen	Human CD95 transfected L cells
Conjugation	Biotin
Alternate Names	CD95; Apoptosis-mediating surface antigen FAS; FAS1; Tumor necrosis factor receptor superfamily member 6; ALPS1A; APT1; FASTM; CD antigen CD95; APO-1; TNFRSF6; FASLG receptor; Apo-1 antigen

### Application Instructions

Application table	Application	Dilution
	ELISA	Assay-dependent
	FACS	10 µl/10 <sup>6</sup> cells
	ICC/IF	Assay-dependent
	IHC-Fr	Assay-dependent
	IHC-P	Assay-dependent

**Application Note** \* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.

### Properties

Form	Liquid
Buffer	PBS and 0.1% Sodium azide.
Preservative	0.1% Sodium azide
Storage instruction	Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note For laboratory research only, not for drug, diagnostic or other use.

## Bioinformation

---

Database links	<a href="#">GeneID: 355 Human</a> <a href="#">Swiss-port # P25445 Human</a>
Gene Symbol	FAS
Gene Full Name	Fas cell surface death receptor
Background	The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor contains a death domain. It has been shown to play a central role in the physiological regulation of programmed cell death, and has been implicated in the pathogenesis of various malignancies and diseases of the immune system. The interaction of this receptor with its ligand allows the formation of a death-inducing signaling complex that includes Fas-associated death domain protein (FADD), caspase 8, and caspase 10. The autoproteolytic processing of the caspases in the complex triggers a downstream caspase cascade, and leads to apoptosis. This receptor has been also shown to activate NF-kappaB, MAPK3/ERK1, and MAPK8/JNK, and is found to be involved in transducing the proliferating signals in normal diploid fibroblast and T cells. Several alternatively spliced transcript variants have been described, some of which are candidates for nonsense-mediated mRNA decay (NMD). The isoforms lacking the transmembrane domain may negatively regulate the apoptosis mediated by the full length isoform. [provided by RefSeq, Mar 2011]
Function	Receptor for TNFSF6/FASLG. The adapter molecule FADD recruits caspase-8 to the activated receptor. The resulting death-inducing signaling complex (DISC) performs caspase-8 proteolytic activation which initiates the subsequent cascade of caspases (aspartate-specific cysteine proteases) mediating apoptosis. FAS-mediated apoptosis may have a role in the induction of peripheral tolerance, in the antigen-stimulated suicide of mature T-cells, or both. The secreted isoforms 2 to 6 block apoptosis (in vitro). [UniProt]
Calculated Mw	38 kDa
PTM	N- and O-glycosylated. O-glycosylated with core 1 or possibly core 8 glycans.