

## ARG22013 anti-CD30 antibody [mCD30.1]

Package: 100 µg  
Store at: -20°C

### Summary

Product Description	Hamster Monoclonal antibody [mCD30.1] recognizes CD30
Tested Reactivity	Ms
Tested Application	BL, FACS
Specificity	Mouse CD30
Host	Hamster
Clonality	Monoclonal
Clone	mCD30.1
Isotype	IgG1
Target Name	CD30
Species	Mouse
Immunogen	Mouse CD30-Mouse IgG1 fusion protein
Conjugation	Un-conjugated
Alternate Names	Tumor necrosis factor receptor superfamily member 8; Ki-1 antigen; CD30; Ki-1; Lymphocyte activation antigen CD30; CD antigen CD30; D1S166E; CD30L receptor

### Application Instructions

Application table	Application	Dilution
	BL	Assay-dependent
	FACS	< 3 µg/10 <sup>6</sup> cells
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

### Properties

Form	Liquid
Buffer	BBS (pH 8.2)
Concentration	0.5 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

### Bioinformation

Database links	<a href="#">GeneID: 21941 Mouse</a> <a href="#">Swiss-port # Q60846 Mouse</a>
Gene Symbol	TNFRSF8
Gene Full Name	tumor necrosis factor receptor superfamily, member 8
Background	CD30 is a member of the TNF-receptor superfamily. This receptor is expressed by activated, but not by resting, T and B cells. TRAF2 and TRAF5 can interact with this receptor, and mediate the signal transduction that leads to the activation of NF-kappaB. This receptor is a positive regulator of apoptosis, and also has been shown to limit the proliferative potential of autoreactive CD8 effector T cells and protect the body against autoimmunity. Two alternatively spliced transcript variants of this gene encoding distinct isoforms have been reported. [provided by RefSeq, Jul 2008]
Function	CD30 is a receptor for TNFSF8/CD30L (PubMed:8391931). May play a role in the regulation of cellular growth and transformation of activated lymphoblasts. Regulates gene expression through activation of NF-kappa-B (PubMed:8999898). [UniProt]
Calculated Mw	64 kDa
PTM	Phosphorylated on serine and tyrosine residues.