

## ARG23422 anti-CD154 / CD40L antibody [TRAP-1]

Package: 100 µg  
Store at: -20°C

### Summary

<b>Product Description</b>	Mouse Monoclonal antibody [TRAP-1] recognizes CD154 / CD40L Mouse anti Human CD154 antibody, clone TRAP-1 recognises the human CD40 ligand, also known as CD154, TNF-related activation protein (TRAP) or T-cell antigen Gp39. CD154 is a 261 amino acid ~ 32kDa single pass, type-1 transmembrane glycoprotein (UniProt: P29965). CD154 is expressed on activated T lymphocytes, predominantly CD4 +ve and also on some basophils and mast cells. Mouse anti Human CD154 antibody, clone TRAP-1 binds to CD154 at an epitope distinct from the CD40 binding site (Krocze et al. 1994).
<b>Tested Reactivity</b>	Hu
<b>Tested Application</b>	FACS
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Clone</b>	TRAP-1
<b>Isotype</b>	IgG1
<b>Target Name</b>	CD154 / CD40L
<b>Species</b>	Human
<b>Immunogen</b>	Mouse myeloma cell line transfected with human CD40L (CD154)
<b>Conjugation</b>	Un-conjugated
<b>Alternate Names</b>	TNFSF5; IMD3; T-cell antigen Gp39; HIGM1; CD40-L; gp39; CD40 ligand; Tumor necrosis factor ligand superfamily member 5; CD40L; CD154; TRAP; CD antigen CD154; hCD40L; IGM; T-BAM; TNF-related activation protein

### Application Instructions

<b>Application table</b>	<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 50%;">Application</th> <th style="width: 50%;">Dilution</th> </tr> </thead> <tbody> <tr> <td>FACS</td> <td>1:10 - 1:50</td> </tr> </tbody> </table>	Application	Dilution	FACS	1:10 - 1:50
Application	Dilution				
FACS	1:10 - 1:50				
<b>Application Note</b>	FACS: Use 10 µl of the suggested working dilution to label 10 <sup>6</sup> cells in 100 µl. * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.				

### Properties

<b>Form</b>	Liquid
<b>Purification</b>	Purification with Protein A.
<b>Buffer</b>	PBS, 0.09% Sodium azide and 0.1% BSA.
<b>Preservative</b>	0.09% Sodium azide
<b>Stabilizer</b>	0.1% BSA

Concentration	1 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

## Bioinformation

---

Gene Symbol	CD40LG
Gene Full Name	CD40 ligand
Background	The protein encoded by this gene is expressed on the surface of T cells. It regulates B cell function by engaging CD40 on the B cell surface. A defect in this gene results in an inability to undergo immunoglobulin class switch and is associated with hyper-IgM syndrome. [provided by RefSeq, Jul 2008]
Function	Mediates B-cell proliferation in the absence of co-stimulus as well as IgE production in the presence of IL-4. Involved in immunoglobulin class switching.  Release of soluble CD40L from platelets is partially regulated by GP IIb/IIIa, actin polymerization, and an matrix metalloproteinases (MMP) inhibitor-sensitive pathway. [UniProt]
Calculated Mw	29 kDa
PTM	The soluble form derives from the membrane form by proteolytic processing.  N-linked glycan is a mixture of high mannose and complex type. Glycan structure does not influence binding affinity to CD40.  Not O-glycosylated. [UniProt]