

ARG23616 anti-CD137L / TNFSF9 antibody [AT113-2]

Package: 100 µg
Store at: -20°C

Summary

Product Description	Rat Monoclonal antibody [AT113-2] recognizes CD137L / TNFSF9
Tested Reactivity	Ms
Tested Application	FACS, IP
Host	Rat
Clonality	Monoclonal
Clone	AT113-2
Isotype	IgG1
Target Name	CD137L / TNFSF9
Species	Mouse
Immunogen	CD137L-Fc Fusion Protein.
Conjugation	Un-conjugated
Alternate Names	4-1BBL; 4-1BB ligand; Tumor necrosis factor ligand superfamily member 9; 4-1BB-L; CD137L

Application Instructions

Application table	Application	Dilution
	FACS	Neat
	IP	Assay-dependent

Application Note FACS: Use 10 µl of the suggested working dilution to label 10⁶ cells in 100 µl.
* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.

Properties

Form	Liquid
Purification	Purification with Protein G.
Buffer	PBS and 0.09% Sodium azide.
Preservative	0.09% Sodium azide
Concentration	1 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol	TNFSF9
Gene Full Name	tumor necrosis factor (ligand) superfamily, member 9
Background	<p>The protein encoded by this gene is a cytokine that belongs to the tumor necrosis factor (TNF) ligand family. This transmembrane cytokine is a bidirectional signal transducer that acts as a ligand for TNFRSF9/4-1BB, which is a costimulatory receptor molecule in T lymphocytes. This cytokine and its receptor are involved in the antigen presentation process and in the generation of cytotoxic T cells. The receptor TNFRSF9/4-1BB is absent from resting T lymphocytes but rapidly expressed upon antigenic stimulation. The ligand encoded by this gene, TNFSF9/4-1BBL, has been shown to reactivate anergic T lymphocytes in addition to promoting T lymphocyte proliferation. This cytokine has also been shown to be required for the optimal CD8 responses in CD8 T cells. This cytokine is expressed in carcinoma cell lines, and is thought to be involved in T cell-tumor cell interaction.[provided by RefSeq, Oct 2008]</p>
Function	<p>Cytokine that binds to TNFRSF9. Induces the proliferation of activated peripheral blood T-cells. May have a role in activation-induced cell death (AICD). May play a role in cognate interactions between T-cells and B-cells/macrophages. [UniProt]</p>
Calculated Mw	27 kDa