

ARG23821 Goat anti-Rat IgG (H+L) antibody (Texas Red) , pre-adsorbed

Package: 500 µg
Store at: 4°C

Summary

Product Description	Texas Red-conjugated Goat Polyclonal antibody recognizes Rat IgG (H+L)
Tested Reactivity	Rat
Tested Application	ELISA, FACS, FLISA, ICC/IF, IHC-Fr, IHC-P, WB
Specificity	Reacts with the heavy and light chains of rat IgG and with the light chains of rat IgM as demonstrated by ELISA, FLISA, and/or flow cytometry; minimal cross reactivity with mouse immunoglobulins
Host	Goat
Clonality	Polyclonal
Isotype	IgG
Target Name	IgG (H+L)
Species	Rat
Conjugation	Texas Red

Application Instructions

Pre Adsorbed	Mouse immunoglobulins and pooled sera.																
Application table	<table><thead><tr><th>Application</th><th>Dilution</th></tr></thead><tbody><tr><td>ELISA</td><td>Assay-dependent</td></tr><tr><td>FACS</td><td>Assay-dependent</td></tr><tr><td>FLISA</td><td>1:100 - 1:400</td></tr><tr><td>ICC/IF</td><td>Assay-dependent</td></tr><tr><td>IHC-Fr</td><td>Assay-dependent</td></tr><tr><td>IHC-P</td><td>Assay-dependent</td></tr><tr><td>WB</td><td>Assay-dependent</td></tr></tbody></table>	Application	Dilution	ELISA	Assay-dependent	FACS	Assay-dependent	FLISA	1:100 - 1:400	ICC/IF	Assay-dependent	IHC-Fr	Assay-dependent	IHC-P	Assay-dependent	WB	Assay-dependent
Application	Dilution																
ELISA	Assay-dependent																
FACS	Assay-dependent																
FLISA	1:100 - 1:400																
ICC/IF	Assay-dependent																
IHC-Fr	Assay-dependent																
IHC-P	Assay-dependent																
WB	Assay-dependent																
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.																

Properties

Form	Liquid
Purification	Affinity purification with immunogen.
Buffer	PBS and 0.1% Sodium azide.
Preservative	0.1% Sodium azide
Concentration	1 mg/ml

Storage instruction Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note For laboratory research only, not for drug, diagnostic or other use.