

## ARG24004 anti-Collagen IV antibody

Package: 50 µl  
Store at: -20°C

### Summary

Product Description	Rabbit Polyclonal antibody recognizes Collagen IV
Tested Reactivity	Bov
Tested Application	ELISA, ICC/IF, IHC-P
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	Collagen IV
Species	Bovine
Immunogen	Type IV collagen extracted from Bovine lens capsules.
Conjugation	Un-conjugated
Alternate Names	BSVD; RATOR; Collagen alpha-1(IV) chain

### Application Instructions

Application table	Application	Dilution
	ELISA	1:2000
	ICC/IF	1:40
	IHC-P	1:500

**Application Note** IHC-P: Antigen Retrieval: treat sections with 0.5% hyaluronidase at RT for 1hr before staining.  
\* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.

### Properties

Form	Liquid
Purification	Purified.
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.


### Bioinformation

Gene Symbol	COL4A1
Gene Full Name	collagen, type IV, alpha 1

<b>Background</b>	This gene encodes a type IV collagen alpha protein. Type IV collagen proteins are integral components of basement membranes. This gene shares a bidirectional promoter with a paralogous gene on the opposite strand. The protein consists of an amino-terminal 7S domain, a triple-helix forming collagenous domain, and a carboxy-terminal non-collagenous domain. It functions as part of a heterotrimer and interacts with other extracellular matrix components such as perlecan, proteoglycans, and laminins. In addition, proteolytic cleavage of the non-collagenous carboxy-terminal domain results in a biologically active fragment known as arresten, which has anti-angiogenic and tumor suppressor properties. Mutations in this gene cause porencephaly, cerebrovascular disease, and renal and muscular defects. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Dec 2014]
<b>Function</b>	Type IV collagen is the major structural component of glomerular basement membranes (GBM), forming a 'chicken-wire' meshwork together with laminins, proteoglycans and entactin/nidogen.  Arresten, comprising the C-terminal NC1 domain, inhibits angiogenesis and tumor formation. The C-terminal half is found to possess the anti-angiogenic activity. Specifically inhibits endothelial cell proliferation, migration and tube formation. Inhibits expression of hypoxia-inducible factor 1alpha and ERK1/2 and p38 MAPK activation. Ligand for alpha1/beta1 integrin. [UniProt]
<b>Calculated Mw</b>	161 kDa
<b>PTM</b>	Lysines at the third position of the tripeptide repeating unit (G-X-Y) are hydroxylated in all cases and bind carbohydrates.  Prolines at the third position of the tripeptide repeating unit (G-X-Y) are hydroxylated in some or all of the chains.  Type IV collagens contain numerous cysteine residues which are involved in inter- and intramolecular disulfide bonding. 12 of these, located in the NC1 domain, are conserved in all known type IV collagens.  The trimeric structure of the NC1 domains is stabilized by covalent bonds between Lys and Met residues.  Proteolytic processing produces the C-terminal NC1 peptide, arresten. [UniProt]
<b>Cellular Localization</b>	Secreted, extracellular space, extracellular matrix, basement membrane. [UniProt]

## Images

---



ARG24004 anti-Collagen IV antibody IHC-P image

Immunohistochemistry: Paraffin-embedded Bovine skin tissue stained with ARG24004 anti-Collagen IV antibody.

---