

ARG43354 anti-SARS-CoV-2 NSP10 antibody

Package: 100 µg
Store at: -20°C

Summary


Product Description	Rabbit Polyclonal antibody recognizes SARS-CoV-2 NSP10
Tested Reactivity	Virus
Tested Application	ELISA, IHC-P, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	SARS-CoV-2 NSP10
Species	Virus
Immunogen	A 10-amino acid synthetic peptide within the last 50 amino acids of SARS-CoV-2 NSP10.
Conjugation	Un-conjugated

Application Instructions

Application table	Application	Dilution
	ELISA	Assay-dependent
	IHC-P	0.5 µg/ml
	WB	1 - 4 µg/ml
Application Note	IHC-P: Antigen Retrieval: Heat mediation was performed in Citrate buffer (pH 6.0). * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	


Properties

Form	Liquid
Purification	Affinity purification with immunogen.
Buffer	PBS and 0.02% Sodium azide.
Preservative	0.02% Sodium azide
Concentration	1 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.




ARG43354 anti-SARS-CoV-2 NSP10 antibody IHC-P image

Immunohistochemistry: Formaldehyde-fixed and paraffin-embedded COVID-19 patient lung tissue. Tissue was blocked with 10% serum for 1 hour at RT. Antigen Retrieval: Heat mediation was performed in Citrate buffer (pH 6.0). Samples were stained with ARG43354 anti-SARS-CoV-2 NSP10 antibody at 0.5 µg/ml dilution, overnight at 4°C. Counter stained with Hematoxylin. Strong signal of SARS-COV-2 NSP10 protein was observed in macrophage of COVID-19 patient lung, but not in non-COVID-19 patient lung.



ARG43354 anti-SARS-CoV-2 NSP10 antibody WB image

Western blot: 30 ng of SARS-CoV-2 NSP10 recombinant protein stained with ARG43354 anti-SARS-CoV-2 NSP10 antibody at 1, 2 or 4 µg/ml dilution (left to right), 1 hour incubation at RT in 5% NFDN/TBST.



ARG43354 anti-SARS-CoV-2 NSP10 antibody ELISA image

Direct ELISA: SARS-CoV-2 NSP10 recombinant protein as coating antigen and ARG43354 anti-SARS-CoV-2 NSP10 antibody as the primary antibody. Detection range is from 1 ng/ml to 3000 ng/ml.