

ARG43833 anti-POLD1 antibody

Package: 100 µl
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes POLD1.
Tested Reactivity	Hu, Ms, Rat
Tested Application	ICC/IF, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	DNA Polymerase Delta 1, Catalytic Subunit
Immunogen	Synthetic peptide of human PolD1.
Conjugation	Un-conjugated
Alternate Names	MDPL; DNA polymerase delta catalytic subunit; DNA polymerase subunit delta p125; CRCS10; POLD; CDC2; EC 2.7.7.7

Application Instructions

Application table	Application	Dilution
	ICC/IF	1:50-1:200
	WB	1:500-1:1000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

Properties

Form	Liquid
Purification	Affinity Purified
Buffer	Tris-Glycine (pH 7.4) with 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA
Preservative	0.01% Sodium azide
Stabilizer	40% Glycerol and 0.05% BSA
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation


Gene Symbol	POLD1
-------------	-------

Gene Full Name	polymerase (DNA directed), delta 1, catalytic subunit
Background	This gene encodes the 125-kDa catalytic subunit of DNA polymerase delta. DNA polymerase delta possesses both polymerase and 3' to 5' exonuclease activity and plays a critical role in DNA replication and repair. Alternatively spliced transcript variants have been observed for this gene, and a pseudogene of this gene is located on the long arm of chromosome 6. [provided by RefSeq, Mar 2012]
Function	Possesses two enzymatic activities: DNA synthesis (polymerase) and an exonucleolytic activity that degrades single stranded DNA in the 3'- to 5'-direction. Required with its accessory proteins (proliferating cell nuclear antigen (PCNA) and replication factor C (RFC) or activator 1) for leading strand synthesis. Also involved in completing Okazaki fragments initiated by the DNA polymerase alpha/primase complex. [UniProt]

Images

ARG43833 anti-POLD1 antibody WB image

Western blot: A562, C6 3T3 and HeLa stained with ARG43833 anti-POLD1 antibody.



ARG43833 anti-POLD1 antibody ICC/IF image

Immunofluorescence: K562 were stained with ARG43833 anti-POLD1 antibody.

