

ARG53805 anti-CD261 / TRAIL R1 antibody [DR-4-02] (PE)

Package: 50 µg
Store at: 4°C

Summary

Product Description	PE-conjugated Mouse Monoclonal antibody [DR-4-02] recognizes CD261 / TRAIL R1
Tested Reactivity	Hu
Tested Application	FACS
Specificity	The clone DR-4-02 recognizes TRAIL-R1 (DR4), a human death receptor 4 (468 amino acids) expressed in most human tissues (spleen, peripheral blood leucocytes, thymus) and in a variety of tumour-derived cell lines.
Host	Mouse
Clonality	Monoclonal
Clone	DR-4-02
Isotype	IgG1
Target Name	CD261 / TRAIL R1
Species	Human
Immunogen	Fusion protein containing the extracellular part of TRAIL-R1 and the constant part of the heavy chain of the human IgG1.
Conjugation	PE
Alternate Names	TNF-related apoptosis-inducing ligand receptor 1; CD antigen CD261; TRAILR-1; DR4; Tumor necrosis factor receptor superfamily member 10A; CD261; Death receptor 4; APO2; TRAIL receptor 1; TRAIL-R1; TRAILR1

Application Instructions

Application table	<table><thead><tr><th>Application</th><th>Dilution</th></tr></thead><tbody><tr><td>FACS</td><td>2 - 5 µg/ml</td></tr></tbody></table>	Application	Dilution	FACS	2 - 5 µg/ml
Application	Dilution				
FACS	2 - 5 µg/ml				
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.				

Properties


Form	Liquid
Purification Note	The purified antibody is conjugated with R-Phycoerythrin (PE) under optimum conditions. The conjugate is purified by size-exclusion chromatography.
Buffer	PBS, 15 mM Sodium azide and 0.2% (w/v) high-grade protease free BSA
Preservative	15 mM Sodium azide
Stabilizer	0.2% (w/v) high-grade protease free BSA
Concentration	0.1 mg/ml

Storage instruction	Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Database links	GeneID: 8797 Human Swiss-port # O00220 Human
Gene Symbol	TNFRSF10A
Gene Full Name	tumor necrosis factor receptor superfamily, member 10a
Background	TRAIL-R1 (CD261, DR4) is a type I transmembrane protein, also called TRAIL receptor 1. The ligand for this DR4 death receptor has been identified and termed TRAIL, which is a member of the TNF family. DR4, as many other receptors (Fas, TNFR1, etc.), mediates apoptosis and NF kappaB activation in many cells and tissues. Apoptosis, a programmed cell death, is a operating process in normal cellular differentiation and development of multicellular organisms. Apoptosis is induced by coupled of certain cytokines (TNF family - TNF, Fas ligand) and their death domain containing receptors (TNFR1, Fas receptor).
Function	Receptor for the cytotoxic ligand TNFSF10/TRAIL. The adapter molecule FADD recruits caspase-8 to the activated receptor. The resulting death-inducing signaling complex (DISC) performs caspase-8 proteolytic activation which initiates the subsequent cascade of caspases (aspartate-specific cysteine proteases) mediating apoptosis. Promotes the activation of NF-kappa-B. [UniProt]
Research Area	Cell Biology and Cellular Response antibody; Cell Death antibody; Immune System antibody
Calculated Mw	50 kDa

Images



ARG53805 anti-CD261 / TRAIL R1 antibody [DR-4-02] (PE) FACS image

Flow Cytometry: CD261-transfected HEK293 cells stained with ARG53805 anti-CD261 / TRAIL R1 antibody [DR-4-02] (PE).