

ARG53832 anti-CD34 antibody [QBEND/10] (PE)

Package: 100 tests
Store at: 4°C

Summary

Product Description	PE-conjugated Mouse Monoclonal antibody [QBEND/10] recognizes CD34 (Endothelial Cell Marker)
Tested Reactivity	Hu, NHuPrm
Species Does Not React With	Rat, Bov, Dog, Sheep
Tested Application	FACS
Specificity	The antibody recognizes the CD34 antigen, some cross reactions with vascular associated adventitia and some basement membranes. Clone QBEND/10 immunoprecipitates a glycoprotein with a relative MW 110kD expressed on haematopoietic cells and on the established myeloid leukaemic cell line KG1A. The clone QBEND/10 recognises the Class II epitope on CD34.
Host	Mouse
Clonality	Monoclonal
Clone	QBEND/10
Isotype	IgG1, kappa
Target Name	CD34
Species	Human
Immunogen	Human endothelial vesicles
Conjugation	PE
Alternate Names	Hematopoietic progenitor cell antigen CD34; CD antigen CD34

Application Instructions

Application table	Application	Dilution
	FACS	20 µl / 10 ⁶ cells
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

Properties

Form	Liquid
Purification Note	The purified antibody is conjugated with R-Phycoerythrin (PE) under optimum conditions. The conjugate is purified by size-exclusion chromatography and adjusted for direct use. No reconstitution is necessary.
Buffer	PBS, 15 mM Sodium azide and 0.2% (w/v) high-grade protease free BSA
Preservative	15 mM Sodium azide
Stabilizer	0.2% (w/v) high-grade protease free BSA
Storage instruction	Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid


repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Database links	GeneID: 947 Human Swiss-port # P28906 Human
Gene Symbol	CD34
Gene Full Name	CD34 molecule
Background	CD34 protein may play a role in the attachment of stem cells to the bone marrow extracellular matrix or to stromal cells. This single-pass membrane protein is highly glycosylated and phosphorylated by protein kinase C. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Aug 2011]
Function	CD34 is a possible adhesion molecule with a role in early hematopoiesis by mediating the attachment of stem cells to the bone marrow extracellular matrix or directly to stromal cells. Could act as a scaffold for the attachment of lineage specific glycans, allowing stem cells to bind to lectins expressed by stromal cells or other marrow components. Presents carbohydrate ligands to selectins. [UniProt]
Research Area	Cancer antibody; Cell Biology and Cellular Response antibody; Controls and Markers antibody; Developmental Biology antibody; Immune System antibody; Neuroscience antibody; Pro-B Cell Marker antibody; Endothelial Cell Marker antibody; Angiogenesis Study antibody
Calculated Mw	41 kDa
PTM	Highly glycosylated. Phosphorylated on serine residues by PKC.

Images



ARG53832 anti-CD34 antibody [QBEND/10] (PE) FACS image

Flow Cytometry: Human peripheral blood stained with ARG53832 anti-CD34 antibody [QBEND/10] (PE) and anti-CD45 antibody (APC).