

Summary

ARG53860 anti-CD45R0 antibody [UCHL1] (PE)

Package: 100 tests Store at: 4°C

| Product Description | PE-conjugated Mouse Monoclonal antibody [UCHL1] recognizes CD45R0 |
|---------------------|--|
| Tested Reactivity | Hu |
| Tested Application | FACS |
| Specificity | The clone UCHL1 recognizes CD45R0, a 180 kDa low molecular weight isoform of the leukocyte common antigen (LCA). The antigen is expressed on a subset of memory/activated T cells and on cortical thymocytes. HLDA III; WS Code NL 826 HLDA III; WS Code T 128 HLDA IV; WS Code NL 31 HLDA V; WS Code BP BP460 HLDA V; WS Code T T-081 HLDA V; WS Code T T-CD45.43 |
| Host | Mouse |
| Clonality | Monoclonal |
| Clone | UCHL1 |
| lsotype | lgG2a |
| Target Name | CD45R0 |
| Species | Human |
| Immunogen | Human IL-2 dependent T cells |
| Conjugation | PE |
| Alternate Names | LY5; GP180; Receptor-type tyrosine-protein phosphatase C; CD45; L-CA; CD antigen CD45; Leukocyte common antigen; CD45R; LCA; T200; EC 3.1.3.48; B220 |

Application Instructions

| Application table | Application | Dilution |
|-------------------|---|--|
| | FACS | 20 μl / 10^6 cells |
| Application Note | * The dilutions indicate should be determined | e recommended starting dilutions and the optimal dilutions or concentrations by the scientist. |


Properties

| Form | Liquid |
|-------------------|---|
| Purification Note | The purified antibody is conjugated with R-Phycoerythrin (PE) under optimum conditions. The conjugate is purified by size-exclusion chromatography and adjusted for direct use. No reconstitution is necessary. |
| Buffer | PBS, 15 mM Sodium azide and 0.2% (w/v) high-grade protease free BSA $$ |

| Preservative | 15 mM Sodium azide |
|---------------------|--|
| Stabilizer | 0.2% (w/v) high-grade protease free BSA |
| Storage instruction | Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use. |
| Note | For laboratory research only, not for drug, diagnostic or other use. |

Bioinformation

| Database links | GenelD: 5788 Human |
|----------------|---|
| | Swiss-port # P08575 Human |
| Gene Symbol | PTPRC |
| Gene Full Name | protein tyrosine phosphatase, receptor type, C |
| Background | CD45 is a member of the protein tyrosine phosphatase (PTP) family. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitosis, and oncogenic transformation. This PTP contains an extracellular domain, a single transmembrane segment and two tandem intracytoplasmic catalytic domains, and thus is classified as a receptor type PTP. This PTP has been shown to be an essential regulator of T- and B-cell antigen receptor signaling. It functions through either direct interaction with components of the antigen receptor complexes, or by activating various Src family kinases required for the antigen receptor signaling. This PTP also suppresses JAK kinases, and thus functions as a regulator of cytokine receptor signaling. Alternatively spliced transcripts variants of this gene, which encode distinct isoforms, have been reported. [provided by RefSeq, Jun 2012] |
| Function | CD45: Protein tyrosine-protein phosphatase required for T-cell activation through the antigen receptor. Acts as a positive regulator of T-cell coactivation upon binding to DPP4. The first PTPase domain has enzymatic activity, while the second one seems to affect the substrate specificity of the first one. Upon T-cell activation, recruits and dephosphorylates SKAP1 and FYN. Dephosphorylates LYN, and thereby modulates LYN activity. (Microbial infection) Acts as a receptor for human cytomegalovirus protein UL11 and mediates binding of UL11 to T-cells, leading to reduced induction of tyrosine phosphorylation of multiple signaling proteins upon T-cell receptor stimulation and impaired T-cell proliferation. [UniProt] |
| Research Area | Developmental Biology antibody; Immune System antibody; Neuroscience antibody; Signaling Transduction antibody; Mouse Inflammatory Cell Marker antibody; B Cell Marker antibody |
| Calculated Mw | 147 kDa |
| PTM | Heavily N- and O-glycosylated. |



ARG53860 anti-CD45R0 antibody [UCHL1] (PE) FACS image

Flow Cytometry: Human peripheral blood stained with ARG53860 anti-CD45R0 antibody [UCHL1] (PE) and anti-CD45RA antibody (APC).