

ARG53891
anti-CD69 antibody [FN50] (PE)Package: 100 tests
Store at: 4°C

Summary

| | |
|---------------------|---|
| Product Description | PE-conjugated Mouse Monoclonal antibody [FN50] recognizes CD69 |
| Tested Reactivity | Hu |
| Tested Application | FACS |
| Specificity | The clone FN50 recognizes CD69, an lymphocyte early activation marker. HLDA IV; WS Code A 91 |
| Host | Mouse |
| Clonality | Monoclonal |
| Clone | FN50 |
| Isotype | IgG1 |
| Target Name | CD69 |
| Species | Human |
| Immunogen | anti- μ -stimulated human B lymphocytes |
| Conjugation | PE |
| Alternate Names | GP32/28; Activation inducer molecule; MLR-3; BL-AC/P26; Leukocyte surface antigen Leu-23; AIM; Early activation antigen CD69; Early T-cell activation antigen p60; EA1; CD antigen CD69; CLEC2C; C-type lectin domain family 2 member C |

Application Instructions

| Application table | Application | Dilution |
|-------------------|-------------|---------------------------|
| | FACS | 20 μ l / 10^6 cells |

Application Note * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.

Properties

| | |
|---------------------|--|
| Form | Liquid |
| Purification Note | The purified antibody is conjugated with R-Phycoerythrin (PE) under optimum conditions. The conjugate is purified by size-exclusion chromatography and adjusted for direct use. No reconstitution is necessary. |
| Buffer | PBS, 15 mM Sodium azide and 0.2% (w/v) high-grade protease free BSA |
| Preservative | 15 mM Sodium azide |
| Stabilizer | 0.2% (w/v) high-grade protease free BSA |
| Storage instruction | Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use. |


Note

For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

| | |
|----------------|---|
| Database links | GeneID: 969 Human Swiss-port # Q07108 Human |
| Gene Symbol | CD69 |
| Gene Full Name | CD69 molecule |
| Background | CD69 (C-type lectin domain family 2 C, CLEC2C, also known as AIM) is one of the earliest inducible cell surface molecules acquired during leukocyte activation. This glycoprotein serves as a lectin-type receptor in lymphocytes, NK cells and platelets; it is involved in lymphocyte proliferation. CD69 expression is counteracted on T cells in the AIDS stage of HIV infection, and may be also predictive for clinical response to chemoimmunotherapy. |
| Function | Involved in lymphocyte proliferation and functions as a signal transmitting receptor in lymphocytes, natural killer (NK) cells, and platelets. [UniProt] |
| Research Area | Developmental Biology antibody; Immune System antibody |
| Calculated Mw | 23 kDa |
| PTM | Constitutive Ser/Thr phosphorylation in both mature thymocytes and activated T-lymphocytes. |

Images



ARG53891 anti-CD69 antibody [FN50] (PE) FACS image

Flow Cytometry: Human PHA-activated peripheral blood stained with ARG53891 anti-CD69 antibody [FN50] (PE).