

## Product datasheet

info@arigobio.com

## ARG54779 anti-Histone H3 monomethyl (Arg17) antibody

Package: 100 μl Store at: -20°C

### **Summary**

**Product Description** Rabbit Polyclonal antibody recognizes Histone H3 monomethyl (Arg17)

**Tested Reactivity** Hu, Ms, Rat

**Tested Application** ICC/IF, IHC-P, WB

Host Rabbit

Polyclonal Clonality

Isotype IgG

**Target Name** Histone H3

**Species** Human

Immunogen Synthetic methylated peptide around Arg17 of Human histone H3 (NP\_003526.1)

Conjugation Un-conjugated

**Alternate Names** Histone H3/f; Histone H3/d; Histone H3/b; Histone H3/c; Histone H3/a; Histone H3/l;

Histone H3/j; Histone H3/k; Histone H3/h; H3/A; H3FA; Histone H3/i

#### **Application Instructions**

Application table	Application	Dilution
	ICC/IF	1:50 - 1:200
	IHC-P	1:50 - 1:200
	WB	1:500 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	HeLa	

#### **Properties**

Form	Liquid	
Purification	Affinity purification with immunogen.	
Buffer	PBS (pH 7.3), 0.02% Sodium azide and 50% Glycerol	
Preservative	0.02% Sodium azide	
Stabilizer	50% Glycerol	
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw	

cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

For laboratory research only, not for drug, diagnostic or other use. Note

#### Bioinformation

Database links <u>GeneID: 8356 Human</u>

Swiss-port # P68431 Human

Gene Symbol HIST1H3J

Gene Full Name histone cluster 1, H3j

Background Histones are basic nuclear proteins that are responsible for the nucleosome structure of the

chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene is intronless and encodes a member of the histone H3 family. Transcripts from this gene lack polyA tails but instead contain a palindromic termination element. This gene is found in the small histone gene cluster on chromosome

6p22-p21.3. [provided by RefSeq, Jul 2008]

Research Area Gene Regulation antibody


Calculated Mw 15 kDa

#### **Images**



# ARG54779 anti-Histone H3 monomethyl (Arg17) antibody ICC/IF image $\,$

Immunofluorescence: 293T cells stained with ARG54779 anti-Histone H3 monomethyl (Arg17) antibody.



#### ARG54779 anti-Histone H3 monomethyl (Arg17) antibody WB image

Western blot: HeLa cell lysate and Recombinant Histone H3 protein expressed in E. coli (negative control) stained with ARG54779 anti-Histone H3 monomethyl (Arg17) antibody .