

ARG55536 anti-Collagen IV antibody

Package: 50 µg
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes Collagen Type IV
Tested Reactivity	Hu, Ms, Rat, Mamm
Tested Application	IHC-Fr, IHC-P, WB
Specificity	Recognizes human Collagen Type IV. Typically less than 1% crossreactivity against other types of collagens was detected by ELISA against purified standards. Reacts with most mammalian Type IV collagens and has negligible crossreactivity with Type I, II, III, V or VI collagens. Non-specific crossreactivity with other human serum proteins or non-collagen extracellular matrix proteins is negligible.
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	Collagen IV
Species	Human
Immunogen	Collagen Type IV from Human and Bovine placenta.
Conjugation	Un-conjugated
Alternate Names	BSVD; RATOR; Collagen alpha-1(IV) chain

Application Instructions

Application table	Application	Dilution
	IHC-Fr	1:50 - 1:200
	IHC-P	1:400
	WB	1:5000 - 1:10000
Application Note	IHC-P: Antigen Retrieval: Boil tissue section in 0.01M Sodium citrate buffer (pH 6.0) for 20 min. * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	


Properties

Form	Liquid
Purification	Affinity purification with immunogen.
Buffer	BBS (pH 8.0), 0.005M EDTA and 0.01% Sodium azide
Preservative	0.01% Sodium azide
Stabilizer	0.005M EDTA
Concentration	1 mg/ml

Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.


Bioinformation

Database links	GeneID: 1282 Human GeneID: 12826 Mouse Swiss-port # P02462 Human Swiss-port # P02463 Mouse
Gene Symbol	COL4A1
Gene Full Name	collagen, type IV, alpha 1
Background	Collagen IV proteins are integral components of basement membranes. This gene shares a bidirectional promoter with a paralogous gene on the opposite strand. The protein consists of an amino-terminal 7S domain, a triple-helix forming collagenous domain, and a carboxy-terminal non-collagenous domain. It functions as part of a heterotrimer and interacts with other extracellular matrix components such as perlecan, proteoglycans, and laminins. In addition, proteolytic cleavage of the non-collagenous carboxy-terminal domain results in a biologically active fragment known as arresten, which has anti-angiogenic and tumor suppressor properties. Mutations in this gene cause porencephaly, cerebrovascular disease, and renal and muscular defects. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Dec 2014]
Function	<p>Collagen IV is the major structural component of glomerular basement membranes (GBM), forming a 'chicken-wire' meshwork together with laminins, proteoglycans and entactin/nidogen.</p> <p>Arresten, comprising the C-terminal NC1 domain, inhibits angiogenesis and tumor formation. The C-terminal half is found to possess the anti-angiogenic activity. Specifically inhibits endothelial cell proliferation, migration and tube formation. Inhibits expression of hypoxia-inducible factor 1alpha and ERK1/2 and p38 MAPK activation. Ligand for alpha1/beta1 integrin. [UniProt]</p>
Research Area	Angiogenesis Study antibody; Basement Membrane Marker antibody
Calculated Mw	161 kDa
PTM	<p>Lysines at the third position of the tripeptide repeating unit (G-X-Y) are hydroxylated in all cases and bind carbohydrates.</p> <p>Prolines at the third position of the tripeptide repeating unit (G-X-Y) are hydroxylated in some or all of the chains.</p> <p>Type IV collagens contain numerous cysteine residues which are involved in inter- and intramolecular disulfide bonding. 12 of these, located in the NC1 domain, are conserved in all known type IV collagens. The trimeric structure of the NC1 domains is stabilized by covalent bonds between Lys and Met residues.</p> <p>Proteolytic processing produces the C-terminal NC1 peptide, arresten.</p>



ARG55536 anti-Collagen IV antibody IHC-P image

Immunohistochemistry: Formalin-fixed and paraffin-embedded Human kidney stained with ARG55536 anti-Collagen IV antibody at 1:400, 45 min at RT. Antigen Retrieval: Boil tissue section in 0.01M Sodium citrate buffer (pH 6.0) for 20 min.



ARG55536 anti-Collagen IV antibody IHC-P image

Immunohistochemistry: Formalin-fixed and paraffin-embedded Human liver stained with ARG55536 anti-Collagen IV antibody at 1:400, 45 min at RT. Antigen Retrieval: Boil tissue section in 0.01M Sodium citrate buffer (pH 6.0) for 20 min.
