

ARG56586 anti-VEGF antibody (Biotin)

Package: 50 µg, 25 µg
Store at: 4°C

Summary

Product Description	Biotin-conjugated Rabbit Polyclonal antibody recognizes VEGF
Tested Reactivity	Hu
Predict Reactivity	Ms, Hm
Tested Application	ELISA, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	VEGF
Species	Human
Immunogen	E. coli derived Recombinant Human VEGF. (APMAEGGGQN HHEVVKFMDV YQRSYCHPIE TLVDIFQEYP DEIEYIFKPS CVPLMRCGGC CNDEGLECVP TEESNITMQI MRIKPHQGQH IGEMSFLQHN KCECRPKKDR ARQENPCGPC SERRKHLFVQ DPQTCKCSCK NTDSRCKARQ LELNERTCRC DKPRR)
Conjugation	Biotin
Alternate Names	MVCD1; Vascular permeability factor; VEGF-A; VPF; VEGF; Vascular endothelial growth factor A

Application Instructions

Application table	Application	Dilution
	ELISA	Direct: 0.25 - 1 µg/ml Sandwich: 0.25 - 1 µg/ml
	WB	0.1 - 0.2 µg/ml
Application Note	Sandwich ELISA (Capture antibody - Detection antibody): ARG56585 (0.5 - 2 µg/ml) - ARG56586 (0.25 - 1 µg/ml)	
	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

Properties


Form	Liquid
Purification	Purified by affinity chromatography.
Buffer	PBS (pH 7.2)
Concentration	1 mg/ml
Storage instruction	Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation


Database links	GeneID: 7422 Human Swiss-port # P15692 Human
Gene Symbol	VEGFA
Gene Full Name	vascular endothelial growth factor A
Background	This gene is a member of the PDGF/VEGF growth factor family and encodes a protein that is often found as a disulfide linked homodimer. This protein is a glycosylated mitogen that specifically acts on endothelial cells and has various effects, including mediating increased vascular permeability, inducing angiogenesis, vasculogenesis and endothelial cell growth, promoting cell migration, and inhibiting apoptosis. Elevated levels of this protein is linked to POEMS syndrome, also known as Crow-Fukase syndrome. Mutations in this gene have been associated with proliferative and nonproliferative diabetic retinopathy. Alternatively spliced transcript variants, encoding either freely secreted or cell-associated isoforms, have been characterized. There is also evidence for the use of non-AUG (CUG) translation initiation sites upstream of, and in-frame with the first AUG, leading to additional isoforms. [provided by RefSeq, Jul 2008]
Function	Growth factor active in angiogenesis, vasculogenesis and endothelial cell growth. Induces endothelial cell proliferation, promotes cell migration, inhibits apoptosis and induces permeabilization of blood vessels. Binds to the FLT1/VEGFR1 and KDR/VEGFR2 receptors, heparan sulfate and heparin. NRP1/Neuropilin-1 binds isoforms VEGF-165 and VEGF-145. Isoform VEGF165B binds to KDR but does not activate downstream signaling pathways, does not activate angiogenesis and inhibits tumor growth. [UniProt]
Highlight	Related Antibody Duos and Panels: ARG30295 VEGF ELISA Antibody Duo Related products: VEGF antibodies ; VEGF ELISA Kits ; VEGF Duos / Panels ; VEGF recombinant proteins ; Anti-Rabbit IgG secondary antibodies ; Related news: The role of HDGF in tumor angiogenesis
Calculated Mw	27 kDa

Images



ARG56586 anti-VEGF antibody (Biotin) standard curve image

Direct ELISA: ARG56586 anti-VEGF antibody (Biotin) at 0.25 - 1.0 $\mu\text{g/ml}$ results of a typical standard run with optical density reading at 405 - 650 nm.



ARG56586 anti-VEGF antibody (Biotin) standard curve image

Sandwich ELISA: ARG56586 anti-VEGF antibody (Biotin) as a detection antibody at 0.25 - 1.0 $\mu\text{g}/\text{ml}$ dilution in combination with [ARG56585](#) anti-VEGF antibody as a capture antibody at 0.5 - 2 $\mu\text{g}/\text{ml}$ dilution. Results of a typical standard run with optical density reading at 405 - 650 nm.