

ARG56630 anti-VEGF antibody

Package: 50 µg
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes VEGF
Tested Reactivity	Ms
Tested Application	ELISA, IHC-P, Neut, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	VEGF
Species	Mouse
Immunogen	E.coli derived Recombinant Mouse VEGF. (MAPTTEGEQK SHEVIKFM DV YQRSYCRPIE TLVDIFQEYP DEIEYIFKPS CVPLMRCAGC CNDEALECVP TSESNITMQI MRIKPHSQSH IGEMSFLQHS RCECRPKKDR TKPEKHCEPC SERRKHLFVQ DPQTCKCSCCK NTDSRCKARQ LELNERTCRC DKPRR)
Conjugation	Un-conjugated
Alternate Names	MVCD1; Vascular permeability factor; VEGF-A; VPF; VEGF; Vascular endothelial growth factor A

Application Instructions

Application table	Application	Dilution
	ELISA	Sandwich: 0.5 - 2.0 µg/ml with ARG56739 as a detection antibody
	IHC-P	2.5 µg/ml
	Neut	0.08 - 0.12 µg/ml (To yield [ND50] of the biological activity of mVEGF (10.00 ng/ml))
	WB	0.1 - 0.2 µg/ml
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

Properties


Form	Liquid
Purification	Affinity purification with immunogen.
Buffer	PBS (pH 7.2)
Concentration	1 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Database links	GeneID: 22339 Mouse Swiss-port # Q00731 Mouse
Gene Symbol	Vegfa
Gene Full Name	vascular endothelial growth factor A
Background	This gene is a member of the PDGF/VEGF growth factor family and encodes a protein that is often found as a disulfide linked homodimer. This protein is a glycosylated mitogen that specifically acts on endothelial cells and has various effects, including mediating increased vascular permeability, inducing angiogenesis, vasculogenesis and endothelial cell growth, promoting cell migration, and inhibiting apoptosis. Elevated levels of this protein is linked to POEMS syndrome, also known as Crow-Fukase syndrome. Mutations in this gene have been associated with proliferative and nonproliferative diabetic retinopathy. Alternatively spliced transcript variants, encoding either freely secreted or cell-associated isoforms, have been characterized. There is also evidence for the use of non-AUG (CUG) translation initiation sites upstream of, and in-frame with the first AUG, leading to additional isoforms. [provided by RefSeq, Jul 2008]
Function	Growth factor active in angiogenesis, vasculogenesis and endothelial cell growth. Induces endothelial cell proliferation, promotes cell migration, inhibits apoptosis and induces permeabilization of blood vessels. Binds to the FLT1/VEGFR1 and KDR/VEGFR2 receptors, heparan sulfate and heparin. NRP1/Neuropilin-1 binds isoforms VEGF-165 and VEGF-145. Isoform VEGF165B binds to KDR but does not activate downstream signaling pathways, does not activate angiogenesis and inhibits tumor growth. [UniProt]
Highlight	Related products: VEGF antibodies; VEGF ELISA Kits; VEGF Duos / Panels; VEGF recombinant proteins; Anti-Rabbit IgG secondary antibodies; Related news: The role of HDGF in tumor angiogenesis
Calculated Mw	27 kDa

Images




ARG56630 anti-VEGF antibody IHC-P image

Immunohistochemistry: Formalin-fixed and paraffin-embedded sections of normal murine kidney. The recommended ARG56630 anti-VEGF antibody concentration is 2.5 µg/ml for one hour at RT. A secondary fluorophore conjugated antibody was used for thirty minutes at RT. Antigen Retrieval: Boil tissue section in a buffer (High pH) followed by cooling at RT for 20 min.


ARG56630 anti-VEGF antibody WB image

Western blot: 250 - 0.24 ng of Mouse VEGF stained with ARG56630 anti-VEGF antibody, under reducing conditions.




ARG56630 anti-VEGF antibody IHC-P image

Immunohistochemistry: Formalin-fixed and paraffin-embedded sections of normal murine kidney. The recommended ARG56630 anti-VEGF antibody concentration is 2.5 $\mu\text{g}/\text{ml}$ for one hour at RT. A secondary fluorophore conjugated antibody was used for thirty minutes at RT. Antigen Retrieval: Boil tissue section in a buffer (High pH) followed by cooling at RT for 20 min.




ARG56630 anti-VEGF antibody IHC-P image


Immunohistochemistry: Formalin-fixed and paraffin-embedded sections of normal murine kidney. The recommended ARG56630 anti-VEGF antibody concentration is 2.5 $\mu\text{g}/\text{ml}$ for one hour at RT. A secondary fluorophore conjugated antibody was used for thirty minutes at RT. Antigen Retrieval: Boil tissue section in a buffer (High pH) followed by cooling at RT for 20 min.



ARG56630 anti-VEGF antibody WB image

Western blot: 250 - 0.24 ng of Mouse VEGF stained with ARG56630 anti-VEGF antibody, under non-reducing conditions.





ARG56630 anti-VEGF antibody standard curve image

Sandwich ELISA: ARG56630 anti-VEGF antibody as a capture antibody at 0.5 - 2.0 $\mu\text{g}/\text{ml}$ combined with ARG56739 anti-VEGF antibody (Biotin) as a detection antibody. Results of a typical standard run with optical density.