

ARG56780
anti-CXCL7 / NAP2 antibody (Biotin)Package: 50 µg
Store at: 4°C

Summary

| | |
|---------------------|--|
| Product Description | Biotin-conjugated Goat Polyclonal antibody recognizes CXCL7 / NAP2 |
| Tested Reactivity | Hu |
| Tested Application | ELISA, WB |
| Host | Goat |
| Clonality | Polyclonal |
| Isotype | IgG |
| Target Name | CXCL7 / NAP2 |
| Species | Human |
| Immunogen | E.coli derived Recombinant Human NAP-2 (CXCL7). (AELRCMCIKT TSGIHPKNIQ SLEVIGKGTH CNQVEVIATL KDGRKICLDP DAPRIKKIVQ KKLAGEDESAD) |
| Conjugation | Biotin |
| Alternate Names | CTAP3; Platelet basic protein; SCYB7; Macrophage-derived growth factor; THBGB; CTAPIII; C-X-C motif chemokine 7; Beta-TG; NAP-2; Small-inducible cytokine B7; TGB; THBGB1; CTAP-III; PBP; MDGF; TC2; Low-affinity platelet factor IV; Leukocyte-derived growth factor; TC1; 74; 73; 1-66; B-TG1; 1-63; CXCL7; TGB1; 1-81; LDGF; LA-PF4 |

Application Instructions

| Application table | Application | Dilution |
|-------------------|-------------|--|
| | ELISA | Direct: 0.25 - 1.0 µg/ml Sandwich: 0.25 - 1.0 µg/ml with ARG56670 as a capture antibody |
| | WB | 0.1 - 0.2 µg/ml |

Application Note * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.

Properties

| | |
|---------------------|--|
| Form | Liquid |
| Purification | Purified by affinity chromatography. |
| Buffer | PBS (pH 7.2) |
| Concentration | 1 mg/ml |
| Storage instruction | Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use. |
| Note | For laboratory research only, not for drug, diagnostic or other use. |

Database links

[GeneID: 5473 Human](#)

[Swiss-port # P02775 Human](#)

Gene Symbol

PPBP

Gene Full Name

pro-platelet basic protein (chemokine (C-X-C motif) ligand 7)

Background

The protein encoded by this gene is a platelet-derived growth factor that belongs to the CXC chemokine family. This growth factor is a potent chemoattractant and activator of neutrophils. It has been shown to stimulate various cellular processes including DNA synthesis, mitosis, glycolysis, intracellular cAMP accumulation, prostaglandin E2 secretion, and synthesis of hyaluronic acid and sulfated glycosaminoglycan. It also stimulates the formation and secretion of plasminogen activator by synovial cells. The protein also is an antimicrobial protein with bactericidal and antifungal activity. [provided by RefSeq, Nov 2014]

Function

LA-PF4 stimulates DNA synthesis, mitosis, glycolysis, intracellular cAMP accumulation, prostaglandin E2 secretion, and synthesis of hyaluronic acid and sulfated glycosaminoglycan. It also stimulates the formation and secretion of plasminogen activator by human synovial cells. NAP-2 is a ligand for CXCR1 and CXCR2, and NAP-2, NAP-2(73), NAP-2(74), NAP-2(1-66), and most potent NAP-2(1-63) are chemoattractants and activators for neutrophils. TC-1 and TC-2 are antibacterial proteins, in vitro released from activated platelet alpha-granules. CTAP-III(1-81) is more potent than CTAP-III desensitize chemokine-induced neutrophil activation. [UniProt]


Calculated Mw

14 kDa

PTM


Proteolytic removal of residues 1-9 produces the active peptide connective tissue-activating peptide III (CTAP-III) (low-affinity platelet factor IV (LA-PF4)).
 Proteolytic removal of residues 1-13 produces the active peptide beta-thromboglobulin, which is released from platelets along with platelet factor 4 and platelet-derived growth factor.
 NAP-2(1-66) is produced by proteolytical processing, probably after secretion by leukocytes other than neutrophils.
 NAP-2(73) and NAP-2(74) seem not be produced by proteolytical processing of secreted precursors but are released in an active form from platelets.

Images



ARG56780 anti-CXCL7 / NAP2 antibody (Biotin) standard curve image

Direct ELISA: ARG56780 anti-CXCL7 / NAP2 antibody (Biotin) at 0.25 - 1.0 µg/ml results of a typical standard run with optical density.



ARG56780 anti-CXCL7 / NAP2 antibody (Biotin) standard curve image

Sandwich ELISA: ARG56780 anti-CXCL7 / NAP2 antibody (Biotin) as a detection antibody at 0.25 - 1.0 µg/ml combined with ARG56670 anti-CXCL7 / NAP-2 antibody as a capture antibody. Results of a typical standard run with optical density.