

ARG56810 anti-Leptin antibody (Biotin)

Package: 50 µg
Store at: 4°C

Summary

Product Description	Biotin-conjugated Rabbit Polyclonal antibody recognizes Leptin
Tested Reactivity	Hu
Tested Application	ELISA, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	Leptin
Species	Human
Immunogen	E.coli derived Recombinant Human Leptin. (MVPIQKVQDD TKTLIKTIPT RINDISHTQS VSSKQKVTGL DFIPGLHPIL TSKMDQTLA VYQQILTSMP SRNVIQISND LENLRDLLHV LAFSKSCHLP WASGLETLD LGGVLEASGY STEVVALSRL QGSLQDMLWQ LDLSPGC)
Conjugation	Biotin
Alternate Names	Leptin; Obese protein; OB; LEPD; Obesity factor; OBS

Application Instructions


Application table	Application	Dilution
	ELISA	Direct: 0.25 - 1.0 µg/ml Sandwich: 0.25 - 1.0 µg/ml with ARG56700 as a capture antibody
	WB	0.1 - 0.2 µg/ml
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

Properties

Form	Liquid
Purification	Purified by affinity chromatography.
Buffer	PBS (pH 7.2)
Concentration	1 mg/ml
Storage instruction	Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.


Database links	GeneID: 3952 Human Swiss-port # P41159 Human
Gene Symbol	LEP
Gene Full Name	leptin
Background	This gene encodes a protein that is secreted by white adipocytes, and which plays a major role in the regulation of body weight. This protein, which acts through the leptin receptor, functions as part of a signaling pathway that can inhibit food intake and/or regulate energy expenditure to maintain constancy of the adipose mass. This protein also has several endocrine functions, and is involved in the regulation of immune and inflammatory responses, hematopoiesis, angiogenesis and wound healing. Mutations in this gene and/or its regulatory regions cause severe obesity, and morbid obesity with hypogonadism. This gene has also been linked to type 2 diabetes mellitus development. [provided by RefSeq, Jul 2008]
Function	May function as part of a signaling pathway that acts to regulate the size of the body fat depot. An increase in the level of LEP may act directly or indirectly on the CNS to inhibit food intake and/or regulate energy expenditure as part of a homeostatic mechanism to maintain constancy of the adipose mass. [UniProt]
Calculated Mw	19 kDa

Images



ARG56810 anti-Leptin antibody (Biotin) standard curve image

Direct ELISA: ARG56810 anti-Leptin antibody (Biotin) at 0.25 - 1.0 μ g/ml results of a typical standard run with optical density.



ARG56810 anti-Leptin antibody (Biotin) standard curve image

Sandwich ELISA: ARG56810 anti-Leptin antibody (Biotin) as a detection antibody at 0.25 - 1.0 μ g/ml combined with ARG56700 anti-Leptin antibody as a capture antibody. Results of a typical standard run with optical density.