

ARG56815 anti-TNFR2 antibody (Biotin)

Package: 50 µg
Store at: 4°C

Summary

Product Description	Biotin-conjugated Rabbit Polyclonal antibody recognizes TNFR2
Tested Reactivity	Hu
Tested Application	ELISA, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	TNFR2
Species	Human
Immunogen	E.coli derived Recombinant Human sTNF Receptor Type II. (MAPEPGSTCR LREYYDQTAQ MCCSKCPGQ HAKVFCTKTS DTVCDSCEDS TYTQLWNWVP ECLSCGSRCS SDQVETQACT REQNRICTCR PGWYCALSQK EGCRLCAPLR KCRPGFGVAR PGTETSDVVC KPCAPGTFSN TTSSTDICRP HQICNVVAIP GNASMDAVCT STSP)
Conjugation	Biotin
Alternate Names	TNFR1B; CD antigen CD120b; p75; CD120b; p75TNFR; TNF-R-II; TNFBR; TBP-II; TNFR80; TBP-2; TNFR2; Tumor necrosis factor receptor 2; Tumor necrosis factor receptor type II; Etanercept; Tumor necrosis factor receptor superfamily member 1B; TNFR-II; TNF-R75; TNF-R2; TNF-RII; p80 TNF-alpha receptor

Application Instructions

Application table	Application	Dilution
	ELISA	Direct: 0.25 - 1.0 µg/ml Sandwich: 0.25 - 1.0 µg/ml with ARG56706 as a capture antibody
	WB	0.1 - 0.2 µg/ml
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

Properties

Form	Liquid
Purification	Purified by affinity chromatography.
Buffer	PBS (pH 7.2)
Concentration	1 mg/ml
Storage instruction	Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation


Database links	GeneID: 7133 Human Swiss-port # P20333 Human
Gene Symbol	TNFRSF1B
Gene Full Name	tumor necrosis factor receptor superfamily, member 1B
Background	The protein encoded by this gene is a member of the TNF-receptor superfamily. This protein and TNF-receptor 1 form a heterocomplex that mediates the recruitment of two anti-apoptotic proteins, c-IAP1 and c-IAP2, which possess E3 ubiquitin ligase activity. The function of IAPs in TNF-receptor signalling is unknown, however, c-IAP1 is thought to potentiate TNF-induced apoptosis by the ubiquitination and degradation of TNF-receptor-associated factor 2, which mediates anti-apoptotic signals. Knockout studies in mice also suggest a role of this protein in protecting neurons from apoptosis by stimulating antioxidative pathways. [provided by RefSeq, Jul 2008]
Function	Receptor with high affinity for TNFSF2/TNF-alpha and approximately 5-fold lower affinity for homotrimeric TNFSF1/lymphotoxin-alpha. The TRAF1/TRAF2 complex recruits the apoptotic suppressors BIRC2 and BIRC3 to TNFRSF1B/TNFR2. This receptor mediates most of the metabolic effects of TNF-alpha. Isoform 2 blocks TNF-alpha-induced apoptosis, which suggests that it regulates TNF-alpha function by antagonizing its biological activity. [UniProt]
Calculated Mw	48 kDa
PTM	Phosphorylated; mainly on serine residues and with a very low level on threonine residues. A soluble form (tumor necrosis factor binding protein 2) is produced from the membrane form by proteolytic processing.

Images



ARG56815 anti-TNFR2 antibody (Biotin) WB image

Western blot: 250 - 0.24 ng of Human sTNF Receptor II stained with ARG56815 anti-TNFR2 antibody (Biotin), under reducing conditions.



ARG56815 anti-TNFR2 antibody (Biotin) WB image

Western blot: 250 - 0.24 ng of Human sTNF Receptor II stained with ARG56815 anti-TNFR2 antibody (Biotin), under non-reducing conditions.