

ARG57700 anti-SPARC antibody

Package: 100 µl
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes SPARC
Tested Reactivity	Hu, Ms, Rat
Tested Application	ICC/IF, IHC-P, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	SPARC
Species	Human
Immunogen	Recombinant protein of Human SPARC.
Conjugation	Un-conjugated
Alternate Names	ON; BM-40; Basement-membrane protein 40; SPARC; Osteonectin; Secreted protein acidic and rich in cysteine

Application Instructions

Application table	Application	Dilution
	ICC/IF	1:50 - 1:200
	IHC-P	1:50 - 1:200
	WB	1:500 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	A375	


Properties

Form	Liquid
Purification	Affinity purified.
Buffer	PBS (pH 7.3), 0.02% Sodium azide and 50% Glycerol.
Preservative	0.02% Sodium azide
Stabilizer	50% Glycerol
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation


Gene Symbol	SPARC
Gene Full Name	secreted protein, acidic, cysteine-rich (osteonectin)
Background	This gene encodes a cysteine-rich acidic matrix-associated protein. The encoded protein is required for the collagen in bone to become calcified but is also involved in extracellular matrix synthesis and promotion of changes to cell shape. The gene product has been associated with tumor suppression but has also been correlated with metastasis based on changes to cell shape which can promote tumor cell invasion. Three transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jun 2015]
Function	Appears to regulate cell growth through interactions with the extracellular matrix and cytokines. Binds calcium and copper, several types of collagen, albumin, thrombospondin, PDGF and cell membranes. There are two calcium binding sites; an acidic domain that binds 5 to 8 Ca(2+) with a low affinity and an EF-hand loop that binds a Ca(2+) ion with a high affinity. [UniProt]
Calculated Mw	35 kDa
Cellular Localization	Secreted, basement membrane, extracellular matrix, extracellular space. [UniProt]

Images



ARG57700 anti-SPARC antibody ICC/IF image

Immunofluorescence: U2OS cells stained with ARG57700 anti-SPARC antibody at 1:100 dilution.



A375

ARG57700 anti-SPARC antibody WB image

Western blot: 25 µg of A375 cell lysate stained with ARG57700 anti-SPARC antibody at 1:3000 dilution.