

ARG66122 anti-SF20 antibody

Package: 50 µg
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes SF20
Tested Reactivity	Ms
Tested Application	ELISA, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	SF20
Species	Mouse
Immunogen	E. coli derived recombinant Mouse SF20. (MVSEPTVPF DVRPGGVVHS FSQDVGPNGK FTCTFTYASQ GGTNEQWQMS LGTSEDSQHF TCTIWRPQ GK SYLYFTQFKA ELRGAEIEYA MAYSKAAFER ESDVPLKSEE FEVTKTAVSH RPGAFKAELS KLVIVAKAAR SEL)
Conjugation	Un-conjugated
Alternate Names	EUROIMAGE1875335; IL27w; MYDGF; Interleukin-25; R33729_1; Myeloid-derived growth factor; IL-25; Stromal cell-derived growth factor SF20; IL25; IL27; SF20; C19orf10

Application Instructions


Application table	Application	Dilution
	ELISA	Sandwich: 0.5 - 2.0 µg/ml with ARG66123 as a detection antibody
	WB	0.1 - 0.2 µg/ml
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

Properties

Form	Liquid
Purification	Affinity purification with immunogen.
Buffer	PBS (pH 7.2)
Concentration	1 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.


Database links	GeneID: 28106 Mouse Swiss-port # Q9CPT4 Mouse
Gene Symbol	Mydgf
Gene Full Name	myeloid derived growth factor
Background	The protein encoded by this gene was previously thought to support proliferation of lymphoid cells and was considered an interleukin. However, this activity has not been reproducible and the function of this protein is currently unknown. [provided by RefSeq, Jul 2008]
Function	Bone marrow-derived monocyte and paracrine-acting protein that promotes cardiac myocyte survival and adaptive angiogenesis for cardiac protection and/or repair after myocardial infarction (MI). Stimulates endothelial cell proliferation through a MAPK1/3-, STAT3- and CCND1-mediated signaling pathway. Inhibits cardiac myocyte apoptosis in a PI3K/AKT-dependent signaling pathway (By similarity). Involved in endothelial cell proliferation and angiogenesis. [UniProt]
Calculated Mw	19 kDa

Images




ARG66122 anti-SF20 antibody WB image

Western blot: 250 - 0.24 ng of Mouse SF-20 stained with ARG66122 anti-SF20 antibody, under non-reducing conditions.



ARG66122 anti-SF20 antibody standard curve image

Sandwich ELISA: ARG66122 anti-SF20 antibody as a capture antibody at 0.5 - 2.0 µg/ml combined with ARG66123 anti-SF20 antibody (Biotin) as a detection antibody. Results of a typical standard run with optical density reading at 405 - 650 nm.



ARG66122 anti-SF20 antibody WB image

Western blot: 250 - 0.24 ng of Mouse SF-20 stained with ARG66122 anti-SF20 antibody, under reducing conditions.