

ARG66144 anti-CD262 / TRAIL R2 antibody

Package: 50 µg
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes CD262 / TRAIL R2
Tested Reactivity	Hu
Tested Application	ELISA, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	CD262 / TRAIL R2
Species	Human
Immunogen	E. coli derived recombinant Human DR5. (MESALITQQD LAPQQRVAPQ QKRSSPSEGL CPPGHHISED GRDCISCKYG QDYSTHWNDL LFCLRCTRCD SGEVELSPCT TTRNTVCQCE EGTFREEDSP EMCRCRTGC PRGMVKVGDG TPWSDIECVH KES)
Conjugation	Un-conjugated
Alternate Names	TRICK2A; TRICK2B; KILLER; TRAILR2; TNF-related apoptosis-inducing ligand receptor 2; DR5; CD antigen CD262; TRICK2; CD262; KILLER/DR5; Tumor necrosis factor receptor superfamily member 10B; Death receptor 5; TRAIL-R2; TRAIL receptor 2; TRICKB; ZTNFR9

Application Instructions

Application table	Application	Dilution
	ELISA	Sandwich: 0.5 - 2.0 µg/ml with ARG66145 as a detection antibody
	WB	0.1 - 0.2 µg/ml
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

Properties

Form	Liquid
Purification	Affinity purification with immunogen.
Buffer	PBS (pH 7.2)
Concentration	1 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformatics

Database links

[GeneID: 8795 Human](#)

[Swiss-port # O14763 Human](#)

Gene Symbol

TNFRSF10B

Gene Full Name

tumor necrosis factor receptor superfamily, member 10b

Background

The protein encoded by this gene is a member of the TNF-receptor superfamily, and contains an intracellular death domain. This receptor can be activated by tumor necrosis factor-related apoptosis inducing ligand (TNFSF10/TRAIL/APO-2L), and transduces an apoptosis signal. Studies with FADD-deficient mice suggested that FADD, a death domain containing adaptor protein, is required for the apoptosis mediated by this protein. Two transcript variants encoding different isoforms and one non-coding transcript have been found for this gene. [provided by RefSeq, Mar 2009]


Function

Receptor for the cytotoxic ligand TNFSF10/TRAIL. The adapter molecule FADD recruits caspase-8 to the activated receptor. The resulting death-inducing signaling complex (DISC) performs caspase-8 proteolytic activation which initiates the subsequent cascade of caspases (aspartate-specific cysteine proteases) mediating apoptosis. Promotes the activation of NF-kappa-B. Essential for ER stress-induced apoptosis. [UniProt]

Calculated Mw


48 kDa

Images



ARG66144 anti-CD262 / TRAIL R2 antibody WB image

Western blot: 250 - 0.24 ng of Human sTRAIL Receptor-2 stained with ARG66144 anti-CD262 / TRAIL R2 antibody, under non-reducing conditions.



ARG66144 anti-CD262 / TRAIL R2 antibody standard curve image

Sandwich ELISA: ARG66144 anti-CD262 / TRAIL R2 antibody as a capture antibody at 0.5 - 2.0 µg/ml combined with ARG66145 anti-DR5 antibody (Biotin) as a detection antibody. Results of a typical standard run with optical density reading at 405 - 650 nm.

ARG66144 anti-CD262 / TRAIL R2 antibody WB image

Western blot: 250 - 0.24 ng of Human sTRAIL Receptor-2 stained with ARG66144 anti-CD262 / TRAIL R2 antibody, under reducing conditions.

