

ARG66145
anti-CD262 / TRAIL R2 antibody (Biotin)Package: 50 µg
Store at: 4°C

Summary

Product Description	Biotin-conjugated Rabbit Polyclonal antibody recognizes CD262 / TRAIL R2
Tested Reactivity	Hu
Tested Application	ELISA, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	CD262 / TRAIL R2
Species	Human
Immunogen	E. coli derived recombinant Human DR5. (MESALITQQD LAPQQRVAPQ QKRSSPSEGL CPPGHHISED GRDCISCKYG QDYSTHWNDL LFCLRCTRCD SGEVELSPCT TTRNTVCQCE EGTFREEDSP EMCRKCRGTC PRGMVKVGDG TPWSDIECVH KES)
Conjugation	Biotin
Alternate Names	TRICK2A; TRICK2B; KILLER; TRAILR2; TNF-related apoptosis-inducing ligand receptor 2; DR5; CD antigen CD262; TRICK2; CD262; KILLER/DR5; Tumor necrosis factor receptor superfamily member 10B; Death receptor 5; TRAIL-R2; TRAIL receptor 2; TRICKB; ZTNFR9

Application Instructions

Application table	Application	Dilution
	ELISA	Direct: 0.25 - 1.0 µg/ml Sandwich: 0.25 - 1.0 µg/ml with ARG66144 as a capture antibody
	WB	0.1 - 0.2 µg/ml

Application Note * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.


Properties

Form	Liquid
Purification	Purified by affinity chromatography.
Buffer	PBS (pH 7.2)
Concentration	1 mg/ml
Storage instruction	Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformatics


Database links	GeneID: 8795 Human Swiss-port # O14763 Human
Gene Symbol	TNFRSF10B
Gene Full Name	tumor necrosis factor receptor superfamily, member 10b
Background	The protein encoded by this gene is a member of the TNF-receptor superfamily, and contains an intracellular death domain. This receptor can be activated by tumor necrosis factor-related apoptosis inducing ligand (TNFSF10/TRAIL/APO-2L), and transduces an apoptosis signal. Studies with FADD-deficient mice suggested that FADD, a death domain containing adaptor protein, is required for the apoptosis mediated by this protein. Two transcript variants encoding different isoforms and one non-coding transcript have been found for this gene. [provided by RefSeq, Mar 2009]
Function	Receptor for the cytotoxic ligand TNFSF10/TRAIL. The adapter molecule FADD recruits caspase-8 to the activated receptor. The resulting death-inducing signaling complex (DISC) performs caspase-8 proteolytic activation which initiates the subsequent cascade of caspases (aspartate-specific cysteine proteases) mediating apoptosis. Promotes the activation of NF-kappa-B. Essential for ER stress-induced apoptosis. [UniProt]
Calculated Mw	48 kDa

Images




ARG66145 anti-CD262 / TRAIL R2 antibody (Biotin) WB image

Western blot: 250 - 0.24 ng of Human sTRAIL Receptor-2 stained with ARG66145 anti-CD262 / TRAIL R2 antibody (Biotin), under non-reducing conditions.




ARG66145 anti-CD262 / TRAIL R2 antibody (Biotin) standard curve image

Direct ELISA: ARG66145 anti-CD262 / TRAIL R2 antibody (Biotin) at 0.25 - 1.0 µg/ml results of a typical standard run with optical density reading at 405 - 650 nm.



ARG66145 anti-CD262 / TRAIL R2 antibody (Biotin) WB image

Western blot: 250 - 0.24 ng of Human sTRAIL Receptor-2 stained with ARG66145 anti-CD262 / TRAIL R2 antibody (Biotin), under reducing conditions.



ARG66145 anti-CD262 / TRAIL R2 antibody (Biotin) standard curve image

Sandwich ELISA: ARG66145 anti-CD262 / TRAIL R2 antibody (Biotin) as a detection antibody at 0.25 - 1.0 µg/ml combined with ARG66144 anti-DR5 antibody as a capture antibody. Results of a typical standard run with optical density reading at 405 - 650 nm.