

## Product datasheet

info@arigobio.com

# ARG67158 anti-ACO1 / Aconitase 1 antibody

Package: 100 μl Store at: -20°C

#### **Summary**

Host

Product Description Rabbit Polyclonal antibody recognizes ACO1 / Aconitase 1

Rabbit

Tested Reactivity Arabi
Tested Application WB

Clonality Polyclonal

Isotype IgG

Target Name ACO1 / Aconitase 1

Species Plant

Immunogen Synthetic peptide corresponding to N-terminus of arabidopsis thaliana ACO1 / Aconitase 1 protein.

Conjugation Un-conjugated

Alternate Names ACO; Aconitate hydratase 1; Aconitase 1; Citrate hydro-lyase 1

### **Application Instructions**

Application table	Application	Dilution
	WB	1:500 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Observed Size	98 kDa	

### **Properties**

Form Liquid

**Purification** Affinity purification with immunogen.

Buffer 0.42% Potassium phosphate (pH 7.3), 0.87% NaCl, 0.01% Sodium azide and 30% Glycerol.

Preservative 0.01% Sodium azide

Stabilizer 30% Glycerol

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed

before use.

Note For laboratory research only, not for drug, diagnostic or other use.

#### Bioinformation

Gene Symbol ACO1

Gene Full Name aconitase 1

Background Encodes an aconitase that can catalyze the conversion of citrate to isocitrate through a cis-aconitate

intermediate, indicating that it may participate in the TCA cycle and other primary metabolic pathways. The protein is believed to accumulate in the mitochondria and the cytosol. It affects CSD2 (At2g28190 - a superoxide dismutase) transcript levels and may play a role in the response to oxidative stress. This

enzyme can also specifically bind to the 5' UTR of CSD2 in vitro.

Function Catalyzes the isomerization of citrate to isocitrate via cis-aconitate; Contributes to oxidative stress

tolerance. [UniProt]

Highlight Related news:

Antibodies for Arabidopsis thaliana;

Calculated Mw 98 kDa

Cellular Localization Cytoplasm, Mitochondrion. [UniProt]

### **Images**



### ARG67158 anti-ACO1 / Aconitase 1 antibody WB image

Western blot: Arabidopsis thaliana stained with ARG67158 anti-ACO1 / Aconitase 1 antibody.