

Product datasheet

info@arigobio.com

ARG70429
Human CXCL9 / MIG recombinant protein (Active) (His-tagged, N-ter)
Story

Package: 100 μg, 20 μg

Store at: -20°C

Summary

Product Description E. coli expressed, His-tagged (N-ter) Active Human CXCL9 / MIG recombinant protein

Tested Application SDS-PAGE

Target Name CXCL9 / MIG

Species Human

A.A. Sequence Thr21 - Thr125

Expression System E. coli

Activity Active

Activity Note Determined by its ability to chemoattract BaF3 cells transfected with mouse CXCR3. The ED50 for this

effect is $< 0.5 \mu g/mL$.

Alternate Names CXCL9; C-X-C Motif Chemokine Ligand 9; SCYB9; Crg-10; Humig; CMK; MIG

Properties

Form Powder

Purification Note Endotoxin level is less than 0.1 EU/μg of the protein, as determined by the LAL test.

Purity > 98% (by SDS-PAGE)

Buffer PBS (pH 7.4)

Reconstitution It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less

than 200 µg/mL and incubate the stock solution for at least 20 min at room temperature to make sure

the protein is dissolved completely.

Storage instruction For long term, lyophilized protein should be stored at -20°C or -80°C. After reconstitution, aliquot and

store at -20°C or -80°C for up to one month. Storage in frost free freezers is not recommended. Avoid

repeated freeze/thaw cycles. Suggest spin the vial prior to opening.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol CXCL9

Gene Full Name C-X-C Motif Chemokine Ligand 9

Background This antimicrobial gene is part of a chemokine superfamily that encodes secreted proteins involved in

immunoregulatory and inflammatory processes. The protein encoded is thought to be involved in T cell trafficking. The encoded protein binds to C-X-C motif chemokine 3 and is a chemoattractant for


lymphocytes but not for neutrophils

Function Cytokine that affects the growth, movement, or activation state of cells that participate in immune and

inflammatory response. Chemotactic for activated T-cells. Binds to CXCR3.

PTM Disulfide bond

Images



ARG70429 Human CXCL9 / MIG recombinant protein (Active) (Histagged, N-ter) SDS-PAGE image $\,$

SDS-PAGE analysis of ARG70429 Human CXCL9 / MIG recombinant protein (Active) (His-tagged, N-ter)