

# Product datasheet

info@arigobio.com

## ARG80212 Mouse CCL2 / MCP1 ELISA Kit

Package: 96 wells Store at: 4°C

# Component

Cat. No.	Component Name	Package	Тетр
ARG80212-001	Antibody-coated microplate	8 X 12 strips	4°C. Unused strips should be sealed tightly in the air-tight pouch.
ARG80212-002	Standard (Lyophilized)	3 X 2 ng/vial	4°C
ARG80212-003	Standard diluent buffer	20 ml	4°C
ARG80212-004	Antibody conjugate concentrate	400 μΙ	4°C
ARG80212-005	Antibody diluent buffer	16 ml	4°C
ARG80212-006	HRP-Streptavidin concentrate	400 μΙ	4°C (Protect from light)
ARG80212-007	HRP-Streptavidin diluent buffer	16 ml	4°C
ARG80212-008	20X Wash buffer	50 ml	4°C
ARG80212-009	TMB substrate	12ml	4°C (Protect from light)
ARG80212-010	STOP solution	12ml	4°C
ARG80212-011	Plate sealer	4 strips	Room temperature

## Summary

Product Description	ARG80212 Mouse CCL2 / MCP1 ELISA Kit is an Enzyme Immunoassay kit for the quantification of Mouse CCL2 / MCP1 in serum, plasma and cell culture supernatants.	
Tested Reactivity	Ms	
Tested Application	ELISA	
Specificity	No significant cross-reactivity or interference with Mouse C10, CTACK, Eotaxin, KC, MCP-2, MDC, MIP-1 alpha and RANTES.	
Target Name	CCL2 / MCP1	
Conjugation	HRP	
Conjugation Note	Substrate: TMB and read at 450 nm	
Sensitivity	7.8 pg/ml	
Sample Type	Serum, plasma and cell culture supernatants.	
Standard Range 15.6 - 1000 pg/ml		

Sample Volume 100 µl

Precision CV: less than 10%

Alternate Names MCP1; Monocyte chemotactic and activating factor; MCAF; Monocyte chemotactic protein 1; Monocyte

secretory protein JE; HSMCR30; Small-inducible cytokine A2; HC11; SMC-CF; GDCF-2; SCYA2; C-C motif

chemokine 2; Monocyte chemoattractant protein 1; MCP-1

#### **Application Instructions**

Assay Time 4 hours

#### **Properties**

Form 96 well

Storage instruction Store the kit at 2-8°C. Keep microplate wells sealed in a dry bag with desiccants. Do not expose test

reagents to heat, sun or strong light during storage and usage. Please refer to the product user manual

for detail temperatures of the components.

Note For laboratory research only, not for drug, diagnostic or other use.

#### Bioinformation

Database links GeneID: 20296 Mouse

Swiss-port # P10148 Mouse

Gene Symbol Ccl2

Gene Full Name chemokine (C-C motif) ligand 2

Background This gene is one of several cytokine genes clustered on the q-arm of chromosome 17. Cytokines are a

family of secreted proteins involved in immunoregulatory and inflammatory processes. The protein encoded by this gene is structurally related to the CXC subfamily of cytokines. Members of this subfamily are characterized by two cysteines separated by a single amino acid. This cytokine displays chemotactic activity for monocytes and basophils but not for neutrophils or eosinophils. It has been implicated in the pathogenesis of diseases characterized by monocytic infiltrates, like psoriasis, rheumatoid arthritis and atherosclerosis. It binds to chemokine receptors CCR2 and CCR4. [provided by

RefSeq, Jul 2008]

Highlight Related products:

MCP1 antibodies; MCP1 ELISA Kits; MCP1 Duos / Panels;

Related news:

HMGB1 in inflammation Inflammatory Cytokines


<u>Detecting MMPs and their non-ECM substrates</u>

New ELISA data calculation tool: Simplify the ELISA analysis by GainData

Research Area Cancer kit; Cell Biology and Cellular Response kit; Immune System kit; Metabolism kit

PTM Processing at the N-terminus can regulate receptor and target cell selectivity. Deletion of the N-

terminal residue converts it from an activator of basophil to an eosinophil chemoattractant.



## ARG80212 Mouse CCL2 / MCP1 ELISA Kit standard curve image

ARG80212 Mouse CCL2 / MCP1 ELISA Kit results of a typical standard run with optical density reading at 450 nm.