

Product datasheet

info@arigobio.com

ARG81342 Mouse CD279 / PD-1 ELISA Kit

Package: 96 wells Store at: 4°C

Component

Cat. No.	Component Name	Package	Temp
ARG81342-001	Antibody-coated microplate	8 X 12 strips	4°C. Unused strips should be sealed tightly in the air-tight pouch.
ARG81342-002	Standard (Lyophilized)	2 X 1.6 ng/vial	4°C
ARG81342-003	Standard diluent buffer	20 ml	4°C
ARG81342-004	Antibody conjugate concentrate	400 μΙ	4°C
ARG81342-005	Antibody diluent buffer	16 ml	4°C
ARG81342-006	HRP-Streptavidin concentrate	400 μΙ	4°C (Protect from light)
ARG81342-007	HRP-Streptavidin diluent buffer	16 ml	4°C
ARG81342-008	20X Wash buffer	50 ml	4°C
ARG81342-009	TMB substrate	12 ml	4°C (Protect from light)
ARG81342-010	STOP solution	12 ml	4°C
ARG81342-011	Plate sealer	4 strips	Room temperature

Summary

Product DescriptionARG81342 Mouse CD279 / PD-1 ELISA Kit is an Enzyme Immunoassay kit for the quantification of Mouse CD279 / PD-1 in serum, plasma and cell culture supernatant.Tested ReactivityMsTested ApplicationELISATarget NameCD279 / PD-1ConjugationHRPConjugation NoteSubstrate: TMB and read at 450 nm.Sensitivity7.8 pg/mlSample TypeSerum, plasma and cell culture supernatant.Standard Range15.6 - 1000 pg/mlSample Volume100 μlPrecisionCV: less than10%		
Tested Application ELISA Target Name CD279 / PD-1 Conjugation Note Substrate: TMB and read at 450 nm. Sensitivity 7.8 pg/ml Sample Type Serum, plasma and cell culture supernatant. Standard Range 15.6 - 1000 pg/ml Sample Volume 100 µl	Product Description	
Target Name CD279 / PD-1 Conjugation HRP Conjugation Note Substrate: TMB and read at 450 nm. Sensitivity 7.8 pg/ml Sample Type Serum, plasma and cell culture supernatant. Standard Range 15.6 - 1000 pg/ml Sample Volume 100 μl	Tested Reactivity	Ms
ConjugationHRPConjugation NoteSubstrate: TMB and read at 450 nm.Sensitivity7.8 pg/mlSample TypeSerum, plasma and cell culture supernatant.Standard Range15.6 - 1000 pg/mlSample Volume100 μl	Tested Application	ELISA
Conjugation NoteSubstrate: TMB and read at 450 nm.Sensitivity7.8 pg/mlSample TypeSerum, plasma and cell culture supernatant.Standard Range15.6 - 1000 pg/mlSample Volume100 μl	Target Name	CD279 / PD-1
Sensitivity 7.8 pg/ml Sample Type Serum, plasma and cell culture supernatant. Standard Range 15.6 - 1000 pg/ml Sample Volume 100 μl	Conjugation	HRP
Sample Type Serum, plasma and cell culture supernatant. Standard Range 15.6 - 1000 pg/ml Sample Volume 100 μl	Conjugation Note	Substrate: TMB and read at 450 nm.
Standard Range 15.6 - 1000 pg/ml Sample Volume 100 μl	Sensitivity	7.8 pg/ml
Sample Volume 100 μl	Sample Type	Serum, plasma and cell culture supernatant.
	Standard Range	15.6 - 1000 pg/ml
Precision CV: less than10%	Sample Volume	100 μΙ
	Precision	CV: less than10%

protein 1; hPD-1

Application Instructions

Assay Time

~ 3.5 hours

Properties

Form

96 well

Storage instruction

Store the kit at $2-8^{\circ}$ C. Keep microplate wells sealed in a dry bag with desiccants. Do not expose test reagents to heat, sun or strong light during storage and usage. Please refer to the product user manual

for detail temperatures of the components.

Note

For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol

PDCD1

Gene Full Name

programmed cell death 1

Background

CD279 / PD-1 is a cell surface membrane protein of the immunoglobulin superfamily. This protein is expressed in pro-B-cells and is thought to play a role in their differentiation. In mice, expression of this gene is induced in the thymus when anti-CD3 antibodies are injected and large numbers of thymocytes undergo apoptosis. Mice deficient for this gene bred on a BALB/c background developed dilated cardiomyopathy and died from congestive heart failure. These studies suggest that this gene product may also be important in T cell function and contribute to the prevention of autoimmune diseases. [provided by RefSeq, Jul 2008]

Function

CD279 / PD-1 is an inhibitory receptor on antigen activated T-cells. It plays a critical role in induction and maintenance of immune tolerance to self (PubMed:21276005). Delivers inhibitory signals upon binding to ligands CD274/PDCD1L1 and CD273/PDCD1LG2 (PubMed:21276005). Following T-cell receptor (TCR) engagement, PDCD1 associates with CD3-TCR in the immunological synapse and directly inhibits T-cell activation. Suppresses T-cell activation through the recruitment of PTPN11/SHP-2: following ligand-binding, PDCD1 is phosphorylated within the ITSM motif, leading to the recruitment of the protein tyrosine phosphatase PTPN11/SHP-2 that mediates dephosphorylation of key TCR proximal signaling molecules, such as ZAP70, PRKCQ/PKCtheta and CD247/CD3zeta.

The PDCD1-mediated inhibitory pathway is exploited by tumors to attenuate anti-tumor immunity and escape destruction by the immune system, thereby facilitating tumor survival (PubMed:28951311). The interaction with CD274/PDCD1L1 inhibits cytotoxic T lymphocytes (CTLs) effector function (PubMed:28951311). The blockage of the PDCD1-mediated pathway results in the reversal of the exhausted T-cell phenotype and the normalization of the anti-tumor response, providing a rationale for cancer immunotherapy (PubMed:22658127, PubMed:25034862, PubMed:25399552). [UniProt]

Highlight

Related products:

PD-1 antibodies; PD-1 ELISA Kits; PD-1 Duos / Panels;


Related news:

New PD-1 ELISA Kit, excellent for preclinical studies or pharmatheutical development

The best solution for PD-1/PD-L1 research

Examining CTL/NK-mediated cytotoxicity by ELISA

New ELISA data calculation tool: Simplify the ELISA analysis by GainData



ARG81342 Mouse CD279 / PD-1 ELISA Kit standard curve image

ARG81342 Mouse CD279 / PD-1 ELISA Kit results of a typical standard run with optical density reading at 450 nm.