

# ARG81676 Mouse sTNFR2 ELISA Kit

Package: 96 wells Store at: 4°C

# Component

Cat. No.	Component Name	Package	Temp
ARG81676-001	Antibody-coated microplate	8 X 12 strips	4°C. Unused strips should be sealed tightly in the air-tight pouch.
ARG81676-002	Standard	2 X 10 ng/vial	4°C
ARG81676-003	Standard/Sample diluent	30 ml (Ready to use)	4°C
ARG81676-004	Antibody conjugate concentrate (100X)	1 vial (100 μl)	4°C
ARG81676-005	Antibody diluent buffer	12 ml (Ready to use)	4°C
ARG81676-006	HRP-Streptavidin concentrate (100X)	1 vial (100 μl)	4°C
ARG81676-007	HRP-Streptavidin diluent buffer	12 ml (Ready to use)	4°C
ARG81676-008	25X Wash buffer	20 ml	4°C
ARG81676-009	TMB substrate	10 ml (Ready to use)	4°C (Protect from light)
ARG81676-010	STOP solution	10 ml (Ready to use)	4°C
ARG81676-011	Plate sealer	4 strips	Room temperature

### Summary

Product Description	ARG81676 Mouse sTNFR2 ELISA Kit is an Enzyme Immunoassay kit for the quantification of Mouse sTNFR2 in serum and cell culture supernatants.	
Tested Reactivity	Ms	
Tested Application	ELISA	
Specificity	There is no detectable cross-reactivity with other relevant proteins.	
Target Name	sTNFR2	
Conjugation	HRP	
Conjugation Note	Substrate: TMB and read at 450 nm.	
Sensitivity	7.8 pg/ml	
Sample Type	Serum and cell culture supernatants.	
Standard Range	15.6 - 1000 pg/ml	
Sample Volume	100 μΙ	

# **Application Instructions**


Assay Time	~ 5 hours
Properties	
Form	96 well
Storage instruction	Store the kit at 2-8°C. Keep microplate wells sealed in a dry bag with desiccants. Do not expose test reagents to heat, sun or strong light during storage and usage. Please refer to the product user manual for detail temperatures of the components.

For laboratory research only, not for drug, diagnostic or other use.

## Bioinformation

Note

Gene Symbol	TNFRSF1B		
Gene Full Name	tumor necrosis factor receptor superfamily, member 1B		
Background	The protein encoded by this gene is a member of the TNF-receptor superfamily. This protein and TNF- receptor 1 form a heterocomplex that mediates the recruitment of two anti-apoptotic proteins, c-IAP1 and c-IAP2, which possess E3 ubiquitin ligase activity. The function of IAPs in TNF-receptor signalling is unknown, however, c-IAP1 is thought to potentiate TNF-induced apoptosis by the ubiquitination and degradation of TNF-receptor-associated factor 2, which mediates anti-apoptotic signals. Knockout studies in mice also suggest a role of this protein in protecting neurons from apoptosis by stimulating antioxidative pathways. [provided by RefSeq, Jul 2008]		
Function	Receptor with high affinity for TNFSF2/TNF-alpha and approximately 5-fold lower affinity for homotrimeric TNFSF1/lymphotoxin-alpha. The TRAF1/TRAF2 complex recruits the apoptotic suppressors BIRC2 and BIRC3 to TNFRSF1B/TNFR2. This receptor mediates most of the metabolic effects of TNF-alpha. Isoform 2 blocks TNF-alpha-induced apoptosis, which suggests that it regulates TNF-alpha function by antagonizing its biological activity. [UniProt]		
Highlight	Related products: <u>TNFR2 antibodies; TNFR2 ELISA Kits;</u> New ELISA data calculation tool: <u>Simplify the ELISA analysis by GainData</u>		
РТМ	Phosphorylated; mainly on serine residues and with a very low level on threonine residues.		
	A soluble form (tumor necrosis factor binding protein 2) is produced from the membrane form by proteolytic processing. [UniProt]		



ARG81676 Mouse sTNFR2 ELISA Kit standard curve image

ARG81676 Mouse sTNFR2 ELISA Kit results of a typical standard run with optical density reading at 450 nm.