

# Product datasheet

info@arigobio.com

## ARG82289 Human IL8 (High sensitive) ELISA Kit

Package: 96 wells Store at: 4°C

## **Summary**

Product Description ARG82289 Human IL8 (High sensitive) ELISA Kit is a high sensitive Enzyme Immunoassay kit for the

quantification of Human IL8 in serum, plasma and cell culture supernatants.

Tested Reactivity Hu

Tested Application ELISA

Specificity Not react with IL1, IFN gamma, HGF, G-CSF, GM-CSF, ANG, AR, CNTF, EGF, FGF and GRO.

Target Name IL8

Conjugation HRP

Conjugation Note Substrate: TMB and read at 450 nm.

Sensitivity 0.78 pg/ml

Sample Type Serum, plasma and cell culture supernatants.

Standard Range 1.56 - 100 pg/ml

Sample Volume  $100 \mu l$ 

Precision Intra-Assay CV: less than 8.5%

Inter-Assay CV: less than 8.5%

Alternate Names IL8/NAP1 form IV; GCP/IL-8 protein IV; NAF; T-cell chemotactic factor; 1-77; Ala-IL-8; Interleukin-8; IL-8;

Protein 3-10C; Lymphocyte-derived neutrophil-activating factor; Neutrophil-activating factor; Granulocyte chemotactic protein 1; LYNAP; NAP-1; Monocyte-derived neutrophil chemotactic factor; 6-77; 7-77; C-X-C motif chemokine 8; GCP1; NAP1; Ser-IL-8; 5-77; GCP/IL-8 protein VI; IL8/NAP1 form I; IL8/NAP1 form VI; Monocyte-derived neutrophil-activating peptide; C-X-C motif; 8-77; 9-77; LUCT;

Chemokine; GCP-1; MDNCF-b; MDNCF-c; IL8/NAP1 form V; LECT; IL8/NAP1 form III; GCP/IL-8 protein III;

Neutrophil-activating protein 1; GCP/IL-8 protein II; IL8/NAP1 form II; GCP/IL-8 protein V; MDNCF;

Emoctakin; GCP/IL-8 protein I; MONAP; IL8

## **Application Instructions**

Assay Time ~ 3.5 hours

### **Properties**

Form 96 well

Storage instruction Store the kit at 2-8°C. Keep microplate wells sealed in a dry bag with desiccants. Do not expose test

reagents to heat, sun or strong light during storage and usage. Please refer to the product user manual

for detail temperatures of the components.

Note For laboratory research only, not for drug, diagnostic or other use.

#### Bioinformation

Gene Symbol CXCL8

Gene Full Name chemokine (C-X-C motif) ligand 8

Background The protein encoded by this gene is a member of the CXC chemokine family. This chemokine is one of

the major mediators of the inflammatory response. This chemokine is secreted by several cell types. It functions as a chemoattractant, and is also a potent angiogenic factor. This gene is believed to play a role in the pathogenesis of bronchiolitis, a common respiratory tract disease caused by viral infection. This gene and other ten members of the CXC chemokine gene family form a chemokine gene cluster in

a region mapped to chromosome 4q. [provided by RefSeq, Jul 2008]

Function IL-8 is a chemotactic factor that attracts neutrophils, basophils, and T-cells, but not monocytes. It is also

involved in neutrophil activation. It is released from several cell types in response to an inflammatory stimulus. IL-8(6-77) has a 5-10-fold higher activity on neutrophil activation, IL-8(5-77) has increased activity on neutrophil activation and IL-8(7-77) has a higher affinity to receptors CXCR1 and CXCR2 as

compared to IL-8(1-77), respectively. [UniProt]

Highlight Related products:

IL8 antibodies; IL8 ELISA Kits; IL8 Duos / Panels; IL8 recombinant proteins;

Related news:

HMGB1 in inflammation
Inflammatory Cytokines
New ELISA data calculation tool:
Simplify the ELISA analysis by GainData

PTM Several N-terminal processed forms are produced by proteolytic cleavage after secretion from at least


peripheral blood monocytes, leukcocytes and endothelial cells. In general, IL-8(1-77) is referred to as

interleukin-8. IL-8(6-77) is the most promiment form.

Citrullination at Arg-27 prevents proteolysis, and dampens tissue inflammation, it also enhances leukocytosis, possibly through impaired chemokine clearance from the blood circulation. [UniProt]

Cellular Localization Secreted. [UniProt]

#### **Images**



ARG82289 Human IL8 (High sensitive) ELISA Kit standard curve image

ARG82289 Human IL8 (High sensitive) ELISA Kit results of a typical standard run with optical density reading at 450 nm.