

## ARG83468 Swine IL8 ELISA Kit

Package: 96 wells

Store at: 4°C

### Summary

Product Description	ARG83468 Swine IL8 ELISA Kit is an Enzyme Immunoassay kit for the quantification of Swine IL8 in Serum, plasma and cell culture supernatants.
Tested Reactivity	Pig
Tested Application	ELISA
Target Name	IL8
Conjugation	HRP
Conjugation Note	Substrate: TMB and read at 450 nm.
Sensitivity	30 pg/ml
Sample Type	Serum, plasma and cell culture supernatants.
Standard Range	62.5 - 4000 pg/ml
Sample Volume	100 µl
Precision	Intra-Assay CV: less than 10% Inter-Assay CV: less than 10%
Alternate Names	IL8/NAP1 form IV; GCP/IL-8 protein IV; NAF; T-cell chemotactic factor; 1-77; Ala-IL-8; Interleukin-8; IL-8; Neutrophil-activating protein 1; GCP/IL-8 protein II; IL8/NAP1 form II; GCP/IL-8 protein V; MDNCF; Protein 3-10C; Lymphocyte-derived neutrophil-activating factor; Neutrophil-activating factor; Granulocyte chemotactic protein 1; LYNAP; NAP-1; Monocyte-derived neutrophil chemotactic factor; 6-77; 7-77; C-X-C motif chemokine 8; GCP1; NAP1; Ser-IL-8; 5-77; GCP/IL-8 protein VI; IL8/NAP1 form I; IL8/NAP1 form VI; Monocyte-derived neutrophil-activating peptide; C-X-C motif; 8-77; 9-77; LUCT; Chemokine; GCP-1; MDNCF-b; MDNCF-c; IL8/NAP1 form V; LECT; IL8/NAP1 form III; GCP/IL-8 protein III; Emotakin; GCP/IL-8 protein I; MONAP; IL8

### Application Instructions

Assay Time	3.5 hours
------------	-----------

### Properties

Form	96 well
Storage instruction	Store the kit at 4°C. Keep microplate wells sealed in a dry bag with desiccants. Do not expose test reagents to heat, sun or strong light during storage and usage. Please refer to the product user manual for detail temperatures of the components.
Note	For laboratory research only, not for drug, diagnostic or other use.

### Bioinformation

Gene Symbol	CXCL8
Gene Full Name	chemokine (C-X-C motif) ligand 8

**Background**

The protein encoded by this gene is a member of the CXC chemokine family. This chemokine is one of the major mediators of the inflammatory response. This chemokine is secreted by several cell types. It functions as a chemoattractant, and is also a potent angiogenic factor. This gene is believed to play a role in the pathogenesis of bronchiolitis, a common respiratory tract disease caused by viral infection. This gene and other ten members of the CXC chemokine gene family form a chemokine gene cluster in a region mapped to chromosome 4q. [provided by RefSeq, Jul 2008]


**Function**

IL-8 is a chemotactic factor that attracts neutrophils, basophils, and T-cells, but not monocytes. It is also involved in neutrophil activation. It is released from several cell types in response to an inflammatory stimulus. IL-8(6-77) has a 5-10-fold higher activity on neutrophil activation, IL-8(5-77) has increased activity on neutrophil activation and IL-8(7-77) has a higher affinity to receptors CXCR1 and CXCR2 as compared to IL-8(1-77), respectively. [UniProt]

**Highlight**

Related products:  
[IL8 antibodies](#); [IL8 ELISA Kits](#); [IL8 Duos / Panels](#); [IL8 recombinant proteins](#); [Anti-Mouse IgG secondary antibodies](#);  
Related news:  
[HMGB1 in inflammation](#)  
[Inflammatory Cytokines](#)

**Images**



ARG83468 Swine IL8 ELISA Kit standard curve image

ARG83468 Swine IL8 ELISA Kit results of a typical standard run with optical density reading at 450 nm.